summary.)

- ÔHIRA, H., 1989 a. Some *Actenicerus*-species of Elateridae from Japan (Coleoptera) (1). *Gekkan-Mushi, Tokyo*, (217): 9–13 (In Japanese with English title.)

- SAITOU, N., & M. NEI, 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. Mol. Biol. Evol., 4: 406-425.
- Swofford, D. L., 2002. PAUP* Phylogenetic Analysis Using Parsimony (*and other methods). Version 4. Sinauer Associates, Sunderland.
- POSADA, D., & K. A. CRANDALL, 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics*, 14: 817–818.
- SAGEGAMI-OBA, R., Y. OBA & H. ÔHIRA, 2007. Phylogenetic relationships of click beetles (Coleoptera: Elateridae) inferred from 28S ribosomal DNA: Insights into the evolution of bioluminescence in Elateridae. *Mol. Phylogenet. Evol.*, **42**: 410–421.
- SIMON, C., F. FRATI, A. BECKENBACH, B. CRESPI, H. LIU & P. FLOOK, 1994. Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Ann. ent. Soc. Am.*, 87: 651-701.
- TARNAWSKI, D., 2001. A world catalogue of Ctenicerini FLEUTIAUX, 1936. Part II (Coleoptera: Elateridae: Athoinae). *Genus, Wrocław*, 12: 277–323.

Elytra, Tokyo, 36(1): 85-86, May 30, 2008

A Host Record of *Thranius variegatus variegatus* (Coleoptera, Cerambycidae)

Shûhei YAMAMOTO

Toho High School, Naka 3-1-10, Kunitachi, Tokyo, 186-0004 Japan

Late in 2007, I recorded *Thranius variegatus variegatus* BATES, 1873 emerged out from dead branches of *Zanthoxylum ailanthoides* SIEBOLD et ZUCC. (Rutaceae) as a distributional record from Kyoto Prefecture, Central Japan (YAMAMOTO, 2007, p. 11). Recentry, I have found out that *Z. ailanthoides* has not been known larval host plant of *T. variegatus variegatus*. In this short paper, I am going to record it as an additional larval host plant of the cerambycid species.

Before going further, I wish to express my hearty thanks to Mr. Satoshi FUJINUMA (Tokyo

Pref.) for giving me invaluable information.

Thranius variegatus variegatus BATES, 1873

[Japanese name: Torafu-hosobane-kamikiri]

Thranius variegatus Bates, 1873, Ann. Mag. nat. Hist., (4), 12: 196; type locality: Nagasaki, Japan. Thranius sapporensis Kano, 1933, Kontyû, Tokyo, 7: 132; type locality: Maruyama, Sapporo City, Hokkaido.

Specimens examined. 1 ♂, 1 ♀, Tsune [崇], Maizuru-shi, northern part of Kyoto Prefecture, Central Japan, 17–III–2006 coll., summer of 2006 emer., S. YAMAMOTO leg. in my coll.; 1 ♀, same collecting data as above, but 10–IV–2007 emer., S. Fujinuma leg. Larval host plants. Mallotus japonicus Müll. Arg. (Euphorbiaceae), Castanopsis spp. (Fagaceae), Ulmus japonica Siebold, Aphananthe aspera Planch (Ulmaceae), Robinia pseudoacacia Linn. (Leguminosae), Zanthoxylum ailanthoides Siebold et Zucc. (Rutaceae: new record.)

Notes. Specimens collected by the author are relatively smaller (body length: 11.3–12.4 mm) than generally known (BL: 13.0–25.5 mm; NIISATO, 2007, p. 464). On the other hand, an individual obtained by Mr. Satoshi FUJINUMA is much larger (BL: 20.0 mm). The hosts were newly dead branches about 6 cm in diameter, with tunnels made by the larvae. In addition to T. variegatus variegatus, ten individuals of Agrilus yamawakii Y. Kurosawa, 1957 also emerged out from the same branches.

References

Kusama, K., & M. Takakuwa, 1984. Cerambycinae. Jpn. Soc. Coleopterol. (ed.), *The Longicorn-Beetles of Japan in Color*, pp. 249–351, pls. 26–48. (In Japanese, with English book title.)

NIISATO, T., 2007. Cerambycinae. *In Ohbayashi*, N., & T. NIISATO (eds.), *Longicorn Beetles of Japan*, pp. 424–512, pls. 22–38. Tokai Univ. Press, Hadano. (In Japanese, with English book title.)

YAMAMOTO, S., 2007. Records of three interesting beetles from Kyoto Prefecture. *Kanagawa-Chûhô*, *Odawara*, (160): 11–12. (In Japanese.)